



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Lack of association of the pregnane X receptor (PXR/NR1I2) gene with inflammatory bowel disease

Citation for published version:

Ho, G-T, Soranzo, N, Tate, SK, Drummond, H, Nimmo, ER, Tenesa, A, Arnott, ID & Satsangi, J 2006, 'Lack of association of the pregnane X receptor (PXR/NR1I2) gene with inflammatory bowel disease: parallel allelic association study and gene wide haplotype analysis', *Gut*, vol. 55, no. 11, pp. 1676-7.
<https://doi.org/10.1136/gut.2006.105106>

Digital Object Identifier (DOI):

[10.1136/gut.2006.105106](https://doi.org/10.1136/gut.2006.105106)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Gut

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



- 9 Lu CC, Sheu BS, Chen TW, *et al*. Host TNF-alpha-1031 and -863 promoter single nucleotide polymorphisms determine the risk of benign ulceration after *H. pylori* infection. *Am J Gastroenterol* 2005;**100**:1274-82.
- 10 Ma SL, Tang NL, Lam LC, *et al*. Association between tumor necrosis factor-alpha promoter polymorphism and Alzheimer's disease. *Neurology* 2004;**62**:307-9.

Lack of association of the pregnane X receptor (PXR/NR1I2) gene with inflammatory bowel disease: parallel allelic association study and gene wide haplotype analysis

The pregnane X receptor gene (PXR/NR1I2) regulates an array of genes involved in the response to xenobiotics.^{1,2} Dysregulation of

this gene may critically influence intestinal barrier defence and susceptibility to inflammatory bowel disease (IBD).³ Recent data from Ireland have suggested strong associations between polymorphisms within the PXR/NR1I2 gene and IBD. Dring *et al* performed a case control study involving 422 patients with IBD (185 ulcerative colitis (UC) and 237 (Crohn's disease (CD)) and 350 healthy controls, using eight candidate polymorphisms in this gene.⁴ Highly significant associations were demonstrated with UC, CD, and IBD as a whole. This effect was most significant for the two individual single nucleotide polymorphisms (SNPs) in the promoter region of this gene; compared between the IBD cohort and controls, rs3814055/-23585 ($p = 0.000008$; odds ratio (OR) 1.62 (95% confidence interval (CI) 1.31-2.00)) and rs1523127/-24381 ($p = 0.0002$; OR 1.50 (95% CI 1.21-1.84)).

We have critically re-evaluated the contribution of these allelic variants of rs1523127/-24381 of the PXR/NR1I2 gene as determinants of disease susceptibility and phenotype in the Scottish population. In addition, we also performed a gene wide association study using a haplotype tagging strategy to assess in detail the overall contribution of this gene to disease susceptibility. A total of 387 UC and 328 CD patients, together with 338 healthy controls (HC), were studied. This study was designed to have 98% power to replicate the previous association with the rs1523127/-24381 variant ($p < 0.05$). In the haplotype analysis, five tagging SNPs (tSNPs) were selected using a multimarker criterion, haplotype $r^2 > 0.80$ to predict all SNPs/haplotypes. This approach was described by Weale and Goldstein and was successfully applied in our previous study of the ABCB1/MDR1 gene.⁵⁻⁸ The exons,

Table 1 Allelic and genotype frequencies of the selected five tagging single nucleotide polymorphisms (tSNPs)

dbSNP ID position	Allele (1/2)	UC	CD	IBD	HC	UC v HC 1 v 2 p value odds ratio 95% CI	UC v HC 1/1 v 2/2 p value odds ratio 95% CI	CD v HC 1 v 2 p value odds ratio 95% CI	CD v HC 1/1 v 2/2 p value odds ratio 95% CI	IBD v HC 1 v 2 p value odds ratio 95% CI	IBD v HC 1/1 v 2/2 p value odds ratio 95% CI
rs1523127	AA	139 (35.9%)	102 (31.1%)	241 (33.7%)	119 (35.6%)	0.96	0.91	0.69	1.00	0.82	1.00
120983729	AG	190 (49.1%)	186 (56.7%)	376 (52.6%)	167 (50.0%)	0.99	0.97	0.95	1.03	0.97	0.99
	GG	58 (15.0%)	40 (12.2%)	98 (13.7%)	48 (14.4%)	0.80-1.22	0.61-1.52	0.76-1.19	0.62-1.69	0.81-1.17	0.66-1.49
	A	468 (60.5%)	390 (59.5%)	858 (60.0%)	405 (60.6%)						
	G	306 (39.5%)	266 (40.5%)	572 (40.0%)	263 (39.4%)						
rs2461823	CC	152 (40.9%)	132 (40.2%)	284 (40.6%)	119 (35.8%)	0.38	0.71	0.46	0.80	0.38	0.66
121002815	CT	175 (47.0%)	157 (47.9%)	332 (47.4%)	174 (52.4%)	1.11	1.11	1.09	1.10	1.09	1.10
	TT	45 (12.1%)	39 (11.9%)	84 (12.0%)	39 (11.7%)	0.88-1.37	0.68-1.81	0.87-1.37	0.66-1.82	0.90-1.32	0.71-1.70
	C	479 (64.4%)	421 (64.2%)	900 (64.3%)	412 (62.0%)						
	T	265 (35.6%)	235 (35.8%)	500 (35.7%)	252 (38.0%)						
	TT	143 (39.7%)	128 (43.2%)	271 (41.3%)	128 (38.5%)	0.79	0.90	0.82	0.62	0.73	0.91
121008187	TC	172 (47.8%)	120 (40.5%)	292 (44.5%)	168 (50.6%)	1.04	1.04	1.04	0.87	1.04	0.96
	CC	45 (12.5%)	48 (16.2%)	93 (14.2%)	42 (12.6%)	0.84-1.29	0.64-1.69	0.82-1.30	0.54-1.42	0.86-1.26	0.63-1.46
	T	458 (63.6%)	376 (63.5%)	834 (63.6%)	424 (62.7%)						
	C	262 (36.4%)	216 (36.5%)	478 (36.4%)	252 (37.3%)						
rs1464603	AA	172 (45.9%)	153 (46.6%)	325 (46.3%)	167 (49.4%)	0.61	1.00	0.72	1.00	0.62	1.00
121009039	AG	159 (42.5%)	138 (42.1%)	297 (42.3%)	130 (38.5%)	0.94	0.98	0.96	1.01	0.95	0.99
	GG	43 (11.5%)	37 (11.3%)	80 (11.4%)	41 (12.1%)	0.75-1.17	0.61-1.58	0.76-1.21	0.62-1.67	0.78-1.15	0.65-1.52
	A	503 (67.2%)	444 (67.7%)	947 (67.5%)	464 (68.6%)						
	G	245 (32.8%)	212 (32.3%)	457 (32.5%)	212 (31.4%)						
rs2472682	TT	42 (11.0%)	35 (11.1%)	77 (11.0%)	39 (11.7%)	0.65	1.00	0.63	0.70	0.97	0.82
121015342	TG	177 (46.2%)	129 (40.8%)	306 (43.8%)	140 (42.2%)	1.06	1.00	0.94	0.90	1.00	0.96
	GG	164 (42.8%)	152 (48.1%)	316 (45.2%)	153 (46.1%)	0.84-1.32	0.62-1.64	0.74-1.19	0.54-1.50	0.82-1.22	0.62-1.47
	T	261 (34.1%)	199 (31.5%)	460 (32.9%)	218 (32.8%)						
	G	505 (65.9%)	433 (68.5%)	938 (67.1%)	446 (67.2%)						

UC, ulcerative colitis; CD, Crohn's disease; HC, healthy controls; 95% CI, 95% confidence interval.

promoter region, and intronic boundaries were resequenced in 16 unrelated Centre d'Etude du Polymorphisme Humain (CEPH) individuals using 10 randomly spaced amplicons across the PXR/NRII2 gene. Haplotypes and their respective frequencies were constructed by further genotyping in 32 CEPH trios.

No association was seen between rs1523127/-24381 ($r^2 = 0.96$ with rs3814055/-23585) SNP and UC, CD, or IBD (A-allelic frequency: 60.5% UC, 59.5% CD, 60.0% IBD, and 60.6% HC; $p = 0.96, 0.69$, and 0.82 , respectively) (table 1). The significant rs6785049 (-7635) variant used in Dring's study was in strong linkage with one of the tSNPs used in this study (rs2472682, $r^2 = 0.86$). In the Scottish dataset, this SNP (rs2472682) was not associated with IBD (T-allele: $p = 0.97$, OR 1.00 (95% CI 0.82-1.22); TT genotype: $p = 0.82$, OR 0.96 (95% CI 0.62-1.47)). Log likelihood analyses comparing overall haplotypic distribution of the five tSNPs demonstrated no associations with UC, CD, or IBD ($p = 0.90, 0.90$, and 0.79 , respectively). There were no associations observed between each of the five tSNPs and the common constructed haplotypes with UC, CD, and IBD, respectively. Genotype-phenotype analyses using the Montreal classification did not show any associations with the studied variants or haplotypes (full data available on request).

These negative data, based on a true candidate gene approach, combine the use of the single marker and multimarker haplotype tagging approach and set a statistical limit to the importance of the contribution of this gene to disease susceptibility in our population. Discordance between the findings of this study and those data presented in Dring's study require clarification and explanation. When the allelic frequencies and homozygosity rates of the rs1523127/-24381 SNP in these two studies are compared, it is of interest to note that differences in the conclusions between these studies may be driven primarily by differences in allelic and genotypic frequencies in healthy controls studied in these datasets (Scottish v Irish controls: A-allele frequencies: 60.6% v 54.8%, respectively, $p = 0.03$, OR 1.27 (95% CI 1.02-1.58); AA-genotype frequencies: 35.6% v 32.2%, respectively, $p = 0.01$, odds ratio 1.75 (95% CI 1.12-2.72)). Differences in frequencies in the Scottish and Irish patient populations were less marked (Scottish v Irish cases: A-allele frequencies: 60.0% v 64.3%, $p = 0.05$, OR 1.20 (95% CI 1.00-1.44); AA-genotype frequencies: 33.7% v 41.1%, $p = 0.17$, odds ratio 1.33 (95% CI 0.90-1.98)). Thus potentially a number of explanations exist, including cryptic population stratification, genetic heterogeneity, or a type 1 error in the initial dataset, each of which may explain the divergent findings in these two populations, in whom previous data had suggested consistent founder effects.⁹ Further studies in these and other Northern European populations may help to resolve this issue formally.

G-T Ho

Molecular Medicine Centre, Gastrointestinal Unit,
University of Edinburgh, Western General Hospital,
Edinburgh, UK

N Soranzo, S K Tate

Department of Clinical and Experimental Epilepsy,
Institute of Neurology, London, UK

H Drummond, E R Nimmo

Molecular Medicine Centre, Gastrointestinal Unit,
University of Edinburgh, Western General Hospital,
Edinburgh, UK

A Tenesa

MRC Human Genetics Unit, Edinburgh, UK

I D Arnott, J Satsangi

Molecular Medicine Centre, Gastrointestinal Unit,
University of Edinburgh, Western General Hospital,
Edinburgh, UK

Correspondence to: Dr G-T Ho, Molecular Medicine
Unit, Gastrointestinal Unit, University of Edinburgh,
Western General Hospital, Edinburgh EH4 2XU, UK;
gwotzerho@aol.com

doi: 10.1136/gut.2006.105106

Dr Gwo-Tzer Ho was supported by the Chief
Scientist's Office, Scottish Executive, UK (2001-04).
Dr Albert Tenesa is funded by Cancer Research UK
grant C348/A3758. Dr Elaine Nimmo and Mrs Hazel
Drummond are supported by a program grant from
the Wellcome Trust, C72789/Z/03/Z.

Conflict of interest: None declared.

References

- Willson TM, Kiewer SA. PXR, CAR and drug metabolism. *Nat Rev Drug Discov* 2002;1:259-66.
- Kiewer SA, Goodwin B, Willson TM. The nuclear pregnane X receptor: a key regulator of xenobiotic metabolism. *Endocr Rev* 2002;23:687-702.
- Langmann T, Moehle C, Maurer R, et al. Loss of detoxification in inflammatory bowel disease: Dysregulation of pregnane X receptor target genes. *Gastroenterology* 2004;127:26-40.
- Dring MM, Goulding CA, Trimble VI, et al. The Pregnane X receptor locus is associated with susceptibility to inflammatory bowel disease. *Gastroenterology* 2006;130:341-8.
- Goldstein DB, Ahmadi KR, Weale ME, et al. Genome scans and candidate gene approaches in the study of common diseases and variable drug responses. *Trends Genet* 2003;19:615-22.
- Weale ME, Depondt C, Macdonald SJ, et al. Selection and evaluation of tagging SNPs in the neuronal-sodium-channel gene SCN1A: implications for linkage-disequilibrium gene mapping. *Am J Hum Genet* 2003;73:551-65.
- Ho GT, Soranzo N, Nimmo ER, et al. ABCB1/MDR1 gene determines susceptibility and phenotype in ulcerative colitis: discrimination of critical variants using a gene-wide haplotype

tagging approach. *Hum Mol Genet* 2006;15:797-805.

- Ahmadi KR, Weale ME, Xue ZY, et al. A single-nucleotide polymorphism tagging set for human drug metabolism and transport. *Nat Genet* 2005;37:84-9.
- Arnott ID, Nimmo ER, Drummond HE, et al. NOD2/CARD15, TLR4 and CD14 mutations in Scottish and Irish Crohn's disease patients: evidence for genetic heterogeneity within Europe? *Genes Immun* 2004;5:417-25.

Genetic association between factor V Leiden and coeliac disease

We identified a family in which a previously unknown genetic association between coeliac disease (CD) and mutation of factor V (factor V Leiden) occurred. The index case was a young woman who abruptly presented with severe ascites due to Budd-Chiari syndrome. At endoscopy, she presented with a scalloped duodenal mucosa suggestive of CD, further confirmed by histology and serological tests. Haematological assessment unveiled the mutation of factor V due to the Arg⁵⁰⁶→Gln mutation at position 506.¹ As shown in fig 1, five siblings were also found to be affected by CD (histology performed in all cases) and were heterozygous carriers of factor V Leiden for the same mutation. The two diseases segregated as one in all cases while no sibling was affected by only one of the two diseases. This pattern of inheritance strongly suggests that the genetic mutation responsible for the onset of CD in this family occurs in a gene which is in high linkage disequilibrium with factor V gene, which is sited on the long arm of chromosome 1. The penetrance of such a gene in our family was found to be 100% as all of the carriers demonstrated coeliac disease at endoscopy.

Although susceptibility to CD is strongly determined by environmental gluten, it is clearly a common genetic disorder. HLA-DQ2 and HLA-DQ8 located in the MHC region on chromosome 6 have been related to a genetic predisposition to CD, although these genes are also present in those not affected by CD.²




Figure 1 Genealogical tree of the family in which an association between factor V Leiden and coeliac disease was unveiled.